

Class of Medication	Generic Name	Brand Name	When to Hold Before Surgery
SGLT-2 Inhibitors Guideline for the Perioperative Management of SGLT2 Inhibitors	Bexagliflozin	Brenzavvy	Hold 72 hours prior to surgery (or will get cancelled)
	Canagliflozin	Invokana	
	Dapagliflozin	Farxiga, Edistride	
	Empagliflozin	Jardiance	
	Ertugliflozin	Steglatro	Hold 96 hours prior to surgery (4 days) (or will get cancelled)
SGLT-2 Inhibitor + metformin	Canagliflozin + metformin	Invokamet	Hold 72 hours prior to surgery (or will get cancelled)
	Dapagliflozin + metformin	Xigduo XR	
	Empagliflozin + metformin	Synjardy/Synjardy XR	
	Ertugliflozin + metformin	Segluromet	Hold 96 hours prior to surgery (or will get cancelled)
SGLT-2 Inhibitor + DPP-4 Inhibitor	Dapagliflozin + saxagliptin	Qtern	Hold 72 hours prior to surgery (or will get cancelled)
	Empagliflozin + linagliptin	Glyxambi	
	Ertugliflozin + sitagliptin	Steglujan	Hold 96 hours prior to surgery (or will get cancelled)
SGLT-2 Inhibitor + metformin + DPP-4 Inhibitor	Empagliflozin + metformin + linagliptin	Trijardy XR	Hold 72 hours prior to surgery (or will get cancelled)
DPP-4 inhibitors	Sitagliptin	Januvia	Hold 24 hours before surgery (or will cancel if taken day of surgery)
	Saxagliptin	Onglyza	
	Linagliptin	Tradjenta	
	Alogliptin	Nesina	
DPP-4 inhibitors + biguanide	Sitagliptin + metformin	Janumet	
	Saxagliptin + metformin	Kombiglyze	
	Linagliptin + metformin	Jentadueto	
	Alogliptin + metformin	Kazano	
DPP-4 inhibitor + thiazolidinedione	Alogliptin + pioglitazone	Oseni	

Class of Medication	Generic Name	Brand Name	When to Hold Before Surgery
GLP-1 receptor agonists (weekly injections)	Dulaglutide	Trulicity	Hold at least 7 days prior to surgery
	Exenatide, extended release	Bydureon	
	Semaglutide	Ozempic (diabetes), Wegovy (weight loss)	
GLP-1 receptor agonists (daily injections or oral meds)	Exenatide, immediate release	Byetta	Hold 24 hours prior to surgery
	Liraglutide	Saxenda (weight loss), Victoza (diabetes)	
	Lixisenatide	Adlyxin	
	Semaglutide	Rybelsus	
GIP/GLP-1 agonist	Tirzepatide	Mounjaro, Zepbound	Hold at least 7 days prior to surgery
Long acting insulins + GLP1 Receptor Agonist	Insulin degludec + liraglutide	Xultophy	Hold 24 hours prior to surgery
	Insulin glargine + lixisenatide	Soliqua	